arXiv:hep-th/9801145v2 3 Feb 1998

hep-th/9801145

String Theory Formulation of anti-de Sitter Black Holes

Danny Birmingham[1]

Department of Mathematical Physics
University College Dublin
Belfield, Dublin 4, Ireland

Abstract

It is shown that the five-dimensional anti-de Sitter black hole is a supersymmetric solution of the low-energy field equations of type IIB string theory compactified on an Einstein space. A statistical interpretation of the mass dependence of the entropy can be obtained from considerations of the three-dimensional BTZ black hole.

January 1998

---

[1] E-mail: dannyb@ollamh.ucd.ie

1 Introduction

The study of black holes in string theory has been the source of recent developments in our understanding of the origin of the Bekenstein-Hawking entropy formula [1]-[3]. In many cases, an important role is played by supersymmetry, and the observation that D-branes carry charge for the RR fields of string theory [4]. In three dimensions, a particularly interesting example of a black hole, known as the BTZ black hole, has been constructed [5, 6], see [7] for a review. The construction is based upon the observation that by performing a quotient of three-dimensional anti-de Sitter space (adS$_3$), one obtains a spacetime with the properties of a black hole. It was observed recently [8] that the microscopic entropy can be understood for certain black holes whose near-horizon geometry is locally equivalent to adS$_3$. This observation is based on the fact that adS$_3$ has an asymptotic symmetry algebra consisting of left and right Virasoro algebras [9, 10]. In particular, the entropy of the extreme BTZ black hole, viewed as a supersymmetric solution of heterotic string theory without the use of RR fields, was computed in [11].

In [12], a higher-dimensional generalization of the BTZ construction was provided. The essential idea is to take a quotient of anti-de Sitter space, yielding a black hole with topology $\mathbf{R}^{d-1} \times S^1$, and an explicit construction in five dimensions was presented. An interesting aspect of these higher-dimensional black holes is that the horizon is a circle, and thus one might hope to be able to understand their entropy from a lower-dimensional point of view. We show that the five-dimensional black hole is a supersymmetric solution of type IIB string theory compactified to adS$_5 \times$K$_5$, where K_5 is a compact internal Einstein space, with only the field strength for the RR 4-form excited. At extremality, the black hole has a non-zero horizon length, proportional to the square root of the mass. A statistical interpretation of this mass dependence can be achieved by relating it to the entropy of the three-dimensional BTZ black hole.

2 Compactification of Type IIB String Theory on adS$_5$

The five-dimensional anti-de Sitter black hole is obtained as a quotient of adS$_5$, to which it is locally equivalent [12]. To obtain this five-dimensional black hole as a supersymmetric solution of type IIB string theory, we seek a compactification to adS$_5\times$K$_5$, where K_5 is a compact internal space. The black hole is then obtained by performing the necessary identifications. We write the 10-dimensional coordinates as $x^M = (x^\mu, y^m)$, with $\mu = 0,1,2,3,4$, and $m = 5,6,7,8,9$, and we follow the conventions of [13]. The covariant field equations for type IIB string theory have been determined in [14]. As observed in [14], the desired compactification can be obtained by setting all the bosonic fields to zero, except for the metric and anti-self-dual 5-form field strength. In this case, the only non-trivial equation of motion is given by

$$R_{MN} = \frac{1}{4^2 \times 6} F_{MN_1N_2N_3N_4} F_N{}^{N_1N_2N_3N_4}, \tag{1}$$

where the 5-form is taken to be anti-self-dual. Thus, we have

$$F_{M_1M_2M_3M_4M_5} = -\frac{1}{5!} \epsilon_{M_1M_2M_3M_4M_5}{}^{N_1N_2N_3N_4N_5} F_{N_1N_2N_3N_4N_5}, \tag{2}$$

with $\epsilon^{0123456789} = 1/\sqrt{-g}$. We wish to obtain a product metric of the form

$$g_{\mu\nu} = g_{\mu\nu}(x), \quad g_{mn} = g_{mn}(y), \quad g_{\mu m} = 0. \tag{3}$$

The required solution follows by choosing the ansatz [14]

$$\begin{aligned} F_{\mu_1\mu_2\mu_3\mu_4\mu_5} &= Q\epsilon_{\mu_1\mu_2\mu_3\mu_4\mu_5}, \\ F_{m_1m_2m_3m_4m_5} &= -Q\epsilon_{m_1m_2m_3m_4m_5}. \end{aligned} \tag{4}$$

It is then straightforward to see that

$$\begin{aligned} R_{\mu\nu} &= -\frac{Q^2}{4}g_{\mu\nu}, \\ R_{mn} &= \frac{Q^2}{4}g_{mn}, \end{aligned} \tag{5}$$

with $R_{\mu m} = 0$, and we note that the 10-dimensional Ricci scalar vanishes.

In order to establish the supersymmetry of the solution, we must show that the Killing spinor equations are satisfied. Namely, we must show that the supersymmetry variations of the fermionic fields vanish in the compactified background. The relevant terms in the supersymmetry transformations take the form [13, 14]

$$\begin{aligned} \delta\psi_M &= \nabla_M\epsilon + \frac{i}{4\times 480}\Gamma^{M_1M_2M_3M_4M_5}\Gamma_M\epsilon F_{M_1M_2M_3M_4M_5} \\ &+ \frac{1}{96}\left(\Gamma_M{}^{M_1M_2M_3} - 9\delta_M{}^{M_1}\Gamma^{M_2M_3}\right)\epsilon^* F_{M_1M_2M_3}, \end{aligned} \tag{6}$$

$$\delta\lambda = i\Gamma^M\epsilon^*\left(\frac{\nabla_M\phi}{1-\phi^*\phi}\right) - \frac{i}{24}\Gamma^{M_1M_2M_3}\epsilon F_{M_1M_2M_3}. \tag{7}$$

A representation of the Dirac matrices which is relevant to the $5+5$ split is given in [15, 16]. The 10-dimensional Dirac matrices are denoted by Γ^A and satisfy

$$\{\Gamma^A, \Gamma^B\} = 2\eta^{AB}, \tag{8}$$

with signature $\eta^{AB} = (-+\cdots+)$. The representation is given by

$$\Gamma^A = (\Gamma^\alpha, \Gamma^a) = (\gamma^\alpha \otimes \mathbf{1}_4, \gamma^6 \otimes \Sigma^a), \tag{9}$$

where the spacetime matrices are 8-dimensional, and the internal matrices are 4-dimensional, satisfying

$$\{\gamma^\alpha, \gamma^\beta\} = 2\eta^{\alpha\beta}, \ \ \{\Sigma^a, \Sigma^b\} = 2\delta^{ab}, \tag{10}$$

with signature $\eta^{\alpha\beta} = (-++++)$. The γ^6 matrix is given explicitly by

$$\gamma^6 = \begin{pmatrix} \mathbf{1}_4 & 0 \\ 0 & -\mathbf{1}_4 \end{pmatrix}, \tag{11}$$

and satisfies

$$\{\gamma^6, \gamma^\alpha\} = 0, \ \ (\gamma^6)^2 = \mathbf{1}_8. \tag{12}$$

The spacetime Dirac matrices can be written as [15, 16]

$$\begin{aligned}
\gamma^0 &= \begin{pmatrix} 0 & i\sigma^1\otimes\mathbf{1}_2 \\ i\sigma^1\otimes\mathbf{1}_2 & 0 \end{pmatrix}, \ \gamma^1 = \begin{pmatrix} 0 & \sigma^3\otimes\mathbf{1}_2 \\ \sigma^3\otimes\mathbf{1}_2 & 0 \end{pmatrix}, \\
\gamma^2 &= \begin{pmatrix} 0 & \sigma^2\otimes\tau^2 \\ \sigma^2\otimes\tau^2 & 0 \end{pmatrix}, \ \gamma^3 = \begin{pmatrix} 0 & \sigma^2\otimes\tau^1 \\ \sigma^2\otimes\tau^1 & 0 \end{pmatrix}, \\
\gamma^4 &= \begin{pmatrix} 0 & \sigma^2\otimes\tau^3 \\ \sigma^2\otimes\tau^3 & 0 \end{pmatrix},
\end{aligned} \tag{13}$$

where σ^i and τ^i are the Pauli matrices. The internal matrices are

$$\begin{aligned}
\Sigma^5 &= \sigma^1\otimes\mathbf{1}_2, \ \Sigma^6 = \sigma^3\otimes\mathbf{1}_2, \ \Sigma^7 = \sigma^2\otimes\tau^2, \\
\Sigma^8 &= \sigma^2\otimes\tau^1, \ \Sigma^9 = \sigma^2\otimes\tau^3,
\end{aligned} \tag{14}$$

with

$$\Gamma^{\alpha\beta} = \gamma^{\alpha\beta}\otimes\mathbf{1}_4, \ \Gamma^{ab} = \mathbf{1}_8\otimes\Sigma^{ab}. \tag{15}$$

A matrix which also enters the analysis is defined by

$$J = \begin{pmatrix} 0 & \mathbf{1}_4 \\ \mathbf{1}_4 & 0 \end{pmatrix}, \tag{16}$$

and satisfies $[J, \gamma^\alpha] = 0$.

We write the 10-dimensional spinor as $\epsilon(x, y) = \eta(x)\otimes\chi(y)$. The Killing spinor equations then become

$$\begin{aligned}
\delta\psi_\mu &= 0 \Rightarrow \ \nabla_\mu\eta = \frac{Q}{8}J\gamma_\mu\eta, \\
\delta\psi_m &= 0 \Rightarrow \ \nabla_m\chi = -i\frac{Q}{8}\Sigma_m\chi,
\end{aligned} \tag{17}$$

with the constraint $iJ\gamma^6\eta = \eta$. The integrability conditions implied by (17) are then precisely the conditions (5). Thus, we have obtained $\mathrm{adS}_5\times\mathrm{K}_5$ as a supersymmetric solution of the low-energy equations of motion of type IIB string theory. The amount of supersymmetry present in five dimensions is then determined by the holonomy of the internal space K_5. One should also note that the constraint on η is consistent with the chirality on ϵ. Defining the 10-dimensional chirality operator as

$$\Gamma^{11} = \Gamma^0\Gamma^1\cdots\Gamma^9, \tag{18}$$

with $(\Gamma^{11})^2 = 1$, we find that

$$\Gamma^{11} = iJ\gamma^6\otimes\mathbf{1}_4. \tag{19}$$

Hence,

$$\Gamma^{11}\epsilon = \epsilon \Rightarrow \ iJ\gamma^6\eta = \eta. \tag{20}$$

3 The Five-Dimensional anti-de Sitter Black Hole

Recently, a higher-dimensional generalization of the BTZ black hole was obtained [12]. The construction is analogous to the three-dimensional case, whereby a certain quotient of anti-de Sitter space is constructed with the properties of a black hole. In particular, the five dimensional case was explicitly constructed. The important point for our purposes is that the topology of the black hole is $\mathbf{R}^{d-1} \times S^1$, with the horizon being given by the S^1 factor. This is to be contrasted with the topology $\mathbf{R}^2 \times S^{d-2}$ of a Schwarzschild black hole. The adS$_5$ black hole is parametrized by two parameters, its mass and angular momentum, and as shown in [12], these can be defined by relating the construction to a Chern-Simons supergravity theory for the supergroup $SU(2,2 \mid N)$ [17].

The line element of the adS$_5$ black hole can be written in the form [12]

$$\begin{aligned} ds^2 &= -\left(\frac{r^2-r_+^2}{\ell^2}\right)\frac{\ell^2}{r_+^2}\cos^2\theta\, dt^2 + \left(\frac{r^2-r_+^2}{\ell^2}\right)^{-1} dr^2 + r^2\frac{\ell^2}{r_+^2}\, d\phi^2 \\ &+ (r^2-r_+^2)\frac{\ell^2}{r_+^2}(d\theta^2+\sin^2\theta\, d\chi^2), \end{aligned} \tag{21}$$

in the range $-\infty < t < \infty, r_+ < r < \infty, 0 < \theta < \pi, 0 \le \chi < 2\pi$. The location of the horizon is specified by $r = r_+$. We note that at the horizon the angular (θ, χ) part of the line element also vanishes. In order to introduce angular momentum, one makes the replacements

$$\begin{aligned} t &\to \frac{r_+}{\ell}t - r_-\phi, \\ \phi &\to \frac{r_-}{\ell^2}t - \frac{r_+}{\ell}\phi, \end{aligned} \tag{22}$$

and identifies points along the new angular coordinate $\phi \sim \phi + 2\pi n$, with $r_- < r_+$. The line element then becomes

$$\begin{aligned} ds^2 &= dt^2\left[\frac{r^2r_-^2}{r_+^2\ell^2} - \left(\frac{r^2-r_+^2}{\ell^2}\right)\cos^2\theta\right] + \left(\frac{r^2-r_+^2}{\ell^2}\right)^{-1} dr^2 + d\phi^2\left[r^2 - \left(\frac{r^2-r_+^2}{\ell^2}\right)\frac{r_-^2\ell^2}{r_+^2}\cos^2\theta\right] \\ &+ dt\, d\phi\left[-\frac{2r^2r_-}{r_+\ell} + \left(\frac{r^2-r_+^2}{\ell^2}\right)\frac{2r_-\ell}{r_+}\cos^2\theta\right] + (r^2-r_+^2)\frac{\ell^2}{r_+^2}(d\theta^2+\sin^2\theta\, d\chi^2). \end{aligned} \tag{23}$$

The Bekenstein-Hawking entropy for the adS$_5$ black hole was computed in [12], and found to be

$$S_{BH} = 4\pi k r_-, \tag{24}$$

where k is the coupling parameter which multiplies the Chern-Simons supergravity action. Since this action is quadratic in curvature, it follows that k has dimensions -1 in length. The mass and angular momentum are given by

$$M = k\frac{2r_+r_-}{\ell^2}, \quad J = k\frac{r_+^2+r_-^2}{\ell}, \tag{25}$$

with the constraint that the horizon exists only for $J \geq M\ell$. The interesting point, as observed in [12], is that the entropy is not proportional to the length of S^1, namely $2\pi r_+$. However, we note that at extremality $J = M\ell$, we have $r_+ = r_-$, and hence the Bekenstein-Hawking entropy is given by

$$S_{BH} = (2\pi r_+)2k. \tag{26}$$

Due to the fact that the horizon for the five-dimensional case is an S^1 factor, one notes that the Bekenstein-Hawking entropy at extremality depends on the square root of the mass, as in the case of the BTZ black hole. For very massive BTZ black holes, the entropy has been understood at the microscopic level in [8, 11]. This understanding is based on the special asymptotic properties of adS$_3$ [9, 10]. Let us recall that in the extreme case, the mass and entropy of the BTZ black hole are given by [5]

$$M_{BTZ} = \frac{r_+^2}{\ell^2}\frac{1}{4G_3}, \quad S_{BTZ} = \frac{2\pi r_+}{4G_3}, \tag{27}$$

where G_3 is the three-dimensional Newton constant with dimensions +1 in length.

In order to relate the mass and entropy of the adS$_5$ black hole to the BTZ case, we need a relationship between the three- and five-dimensional Newton constants. Within string theory, it is natural to identify $k = \alpha'/8G_5$, where G_5 is the five-dimensional Newton constant with dimensions +3 in length. The entropy is then given by

$$S_{BH} = (2\pi r_+)\frac{\alpha'}{4G_5}. \tag{28}$$

However, in order of magnitude, we have $G_3 \sim G_5/\alpha'$. We thus see that the mass of the five-dimensional black hole is of the order of the mass of the BTZ black hole. Upon this identification, one then finds that the entropy of the adS$_5$ black hole is of the order of the BTZ entropy (which has a statistical interpretation [8, 11, 18]). In order words, this identification yields the correct mass dependence of the five-dimensional entropy. In this sense, the analysis is similar in spirit to the correspondence principle presented in [19, 20]. However, further work is required in order to fix the numerical coefficient.

4 Conclusions

The construction of conserved charges for the five-dimensional anti-de Sitter black hole relied on the formulation of a Chern-Simons supergravity theory for the supergroup $SU(2,2 \mid N)$ [17]. In this regard, it is worth remarking that a computation of the entropy for the BTZ black hole from the point of view of three-dimensional Chern-Simons theory with boundary was provided in [18]. The computation is based essentially on the observation that the horizon dynamics on the boundary is controlled by a WZW model. It would be interesting to see if a similar computation in the $SU(2,2 \mid N)$ Chern-Simons theory sheds further light on the entropy. Indeed, there has been recent progress in understanding the connection between singleton field theories in anti-de Sitter space and the D-brane picture of black hole entropy [21]. Further considerations are contained in [22]-[24]. One should also mention that the self-dual 3-brane of type IIB string theory has been discussed in the context of vacuum interpolation between 10-dimensional Minkowski space and adS$_5 \times$ S^5 [25, 26].

Another interesting observation is the fact that the Chern-Simons supergravity theory for the anti-de Sitter group can be constructed up to a maximal dimension of seven [17]. One might therefore expect to be able to construct conserved charges for the seven-dimensional anti-de Sitter black hole, as in the five-dimensional case. Furthermore, the adS_7 black hole could then be obtained as a compactification $\mathrm{adS}_7 \times \mathrm{K}_4$ of 11-dimensional supergravity, with the 4-form field strength being proportional to the volume form of the compact internal space K_4 [27, 28]. In this way, the adS_7 black hole could be given an interpretation as a solution of the low-energy limit of M Theory, with the 2-brane field excited.

Acknowledgements

I would like to thank K. Sfetsos for valuable correspondence on the entropy discussion. This work was supported by Forbairt grant SC/96/603.

References

[1] A. Strominger and C. Vafa, *Microscopic Origin of the Bekenstein-Hawking Entropy*, Phys. Lett. B379 (1996) 99, hep-th/9601029.

[2] C. Callan and J. Maldacena, *D-brane Approach to Black Hole Quantum Mechanics*, Nucl. Phys. B472 (1996) 591, hep-th/9602043.

[3] See, for example, G. Horowitz, *Quantum States of Black Holes*, gr-qc/9704072; J. Maldacena, *Black Holes in String Theory*, hep-th/9607235.

[4] J. Polchinski, *Dirichlet-branes and Ramond-Ramond Charges*, Phys. Rev. Lett. 75 (1995) 4724, hep-th/9510017.

[5] M. Bañados, C. Teitelboim, and J. Zanelli, *Black Hole in Three-Dimensional Spacetime*, Phys. Rev. Lett. 69 (1992) 1849, hep-th/9204099.

[6] M. Bañados, M. Henneaux, C. Teitelboim, and J. Zanelli, *Geometry of the* $2+1$ *Black Hole*, Phys. Rev. D48 (1993) 1506, gr-qc/9302012.

[7] S. Carlip, *The* $(2+1)$*-Dimensional Black Hole*, Class. Quant. Grav. 12 (1995) 2853, gr-qc/9506079.

[8] A. Strominger, *Black Hole Entropy from Near-Horizon Microstates*, hep-th/9712251.

[9] J.D. Brown and M. Henneaux, *Central Charges in the Canonical Realization of Asymptotic Symmetries: An Example from Three Dimensional Gravity*, Commun. Math. Phys. 104 (1986) 207.

[10] O. Coussaert and M. Henneaux, *Supersymmetry of the* $(2+1)$*-Dimensional Black Holes*, Phys. Rev. Lett. 72 (1994) 183, hep-th/9310194.

[11] D. Birmingham, I. Sachs, and S. Sen, *Entropy of Three-Dimensional Black Holes in String Theory*, hep-th/9801019.

[12] M. Bañados, *Constant Curvature Black Holes*, gr-qc/9703040.

[13] M.J. Duff, R.R. Khuri, and J.X. Lu, *String Solitons*, Phys. Rep. 259 (1995) 213, hep-th/9412184.

[14] J.H. Schwarz, *Covariant Field Equations of Chiral* $N = 2$ $D = 10$ *Supergravity*, Nucl. Phys. B226 (1983) 269.

[15] C. Wetterich, *Dimensional Reduction of Weyl, Majorana and Majorana-Weyl Spinors*, Nucl. Phys. B222 (1983) 20.

[16] J.H. Schwarz, in *Superstrings and Supergravity*, Proceedings of the 28th Scottish Universities Summer School in Physics, ed. A.T. Davies and D.G. Sutherland, 1986.

[17] A.H. Chamseddine, *Topological Gravity and Supergravity in Various Dimensions*, Nucl. Phys. B346 (1990) 213.

[18] S. Carlip, *Statistical Mechanics of the* $(2+1)$- *Dimensional Black Hole*, Phys. Rev. D51 (1995) 632, gr-qc/9409052; *The Statistical Mechanics of the Three-Dimensional Euclidean Black Hole*, Phys. Rev. D55 (1997) 878, gr-qc/9606043.

[19] L. Susskind, Some Speculations about Black Hole Entropy in String Theory, hep-th/9309145.

[20] G.T. Horowitz and J. Polchinski, *A Correspondence Principle for Black Holes and Strings*, Phys. Rev. D55 (1997) 6189, hep-th/9612146.

[21] K. Sfetsos and K. Skenderis, *Microscopic Derivation of the Bekenstein-Hawking Entropy Formula for Non-Extremal Black Holes*, hep-th/9711138.

[22] J. Maldacena, *The Large* N *Limit of Superconformal Field Theories and Supergravity*, hep-th/9711200.

[23] S. Ferrara and C. Fronsdal, *Conformal Maxwell Theory as a Singleton Field Theory on* ADS_5*, IIB Three Branes and Duality*, hep-th/9712239.

[24] H.J. Boonstra, B. Peeters, and K. Skenderis, *Branes and anti-de Sitter spacetimes*, hep-th/9801076.

[25] M.J. Duff and J.X. Lu, *The Self-Dual Type IIB Superthreebrane*, Phys. Lett. B273 (1991) 409.

[26] G.W. Gibbons and P.K. Townsend, *Vacuum Interpolation in Supergravity via Super* p*-branes*, Phys. Rev. Lett. 71 (1993) 3754, hep-th/9307049.

[27] P.G.O. Freund and M.A. Rubin, *Dynamics of Dimensional Reduction*, Phys. Lett. 97B (1980) 233.

[28] M.J. Duff, B.E.W. Nilsson, and C.N. Pope, *Kaluza-Klein Supergravity*, Phys. Rep. 130 (1986) 1.